

Phenomenal Characteristics of the Peripheral Drift Illusion

Akiyoshi KITAOKA* and Hiroshi ASHIDA**

* Department of Psychology, College of Letters, Ritsumeikan University

56-1 Toji-in Kitamachi, Kita-ku, Kyoto 603-8577, Japan

** Graduate School of Letters, Kyoto University

Yoshida-honmachi, Sakyo-ku, Kyoto 606-8501, Japan

The peripheral drift illusion refers to an anomalous motion illusion that can be observed in peripheral vision¹⁻³⁾ (Fig. 1). Although this illusion has been characterized by gratings that have sawtooth luminance profiles, we demonstrate that ones with stepwise profiles are more effective. Moreover, the order of four regions of different luminances is critical, i.e. the combination of black and dark-gray or the combination of white and light-gray. Specifically, illusory motion tends to appear in the direction from a black region to an adjacent dark-gray region or in the direction from a white region to an adjacent light-gray region (Fig. 2). This order enhances the peripheral drift




Fig.1 The peripheral drift illusion. When observers see this figure in the peripheral vision, the circle appears to rotate slowly. Although Fraser and Wilcox¹⁾ reported that the direction of illusory rotation was clockwise or counterclockwise depending on individuals, recent studies^{2,3)} mentioned only clockwise rotation in this case.

illusion as shown in Fig. 3.

We point out two novel factors that affect the strength of illusory motion. (1) The peripheral drift illusion is enhanced by stepwise luminance profiles, as mentioned above. Stimuli with smooth luminance profiles, like Fig. 1, give weaker illusion. (2) The peripheral drift illusion is enhanced by fragmented or curved edges. Stimuli made up of long edges give weaker illusion. Fig. 4 is produced




Fig.2 The phenomenal rules of the peripheral drift illusion we here propose, i.e. black to dark-gray and white to light-gray. Arrows indicate assumed local motion signals and their strengths, which originate from black or white areas.




Fig.3 Enhancement of the peripheral drift illusion by our new rules. When observers see this figure in the peripheral vision, the circle appears to rotate clockwise. This illusory rotation is stronger than that in Fig. 1.

according to these rules and shows illusory rotation even stronger than that in Fig. 3. There are a variety of applied figures of the peripheral drift illusion on the first author's webpage: <http://www.ritsumei.ac.jp/~akitaoka/index-e.html>.

Although this study only reveals phenomenological or design rules to obtain the peripheral drift illusion with large illusion magnitude, we are seeking an explanation on the basis of motion detector models⁴⁾. It is also expected that these rules can contribute to the study of motion perception to elucidate anomalous motion illusions⁵⁻¹¹⁾ such as the Ouchi illusion (Fig. 5).

Acknowledgment.

This research was supported by Grant-in-Aid (the




Fig.4 Further enhancement of the peripheral drift illusion by our further rules. When observers see this figure in the peripheral vision, the circle appears to rotate clockwise. This illusory rotation is even stronger than that in Fig. 3.




Fig.5 The Ouchi illusion, the most typical among anomalous motion illusions. The inset appears to move while the surround does not.

Heisei 14th grant entitled "Internet edutainment using anomalous motion illusions") awarded from the Hayao Nakayama Foundation for Science, Technology and Culture, and Grant-in-Aid for Scientific Research (Nos. 14310036 and 14310045) awarded from the Japan Society for the Promotion of Science.

References

- 1) A. Fraser and K. J. Wilcox: Perception of illusory movement. *Nature*, **281**, 565-566, 1979.
- 2) J. Faubert and A. M. Herbert: The peripheral drift illusion: A motion illusion in the visual periphery. *Perception*, **28**, 617-621, 1999.
- 3) G. Naor-Raz and R. Sekuler: Perceptual dimorphism in visual motion from stationary patterns. *Perception*, **29**, 325-335, 2000.
- 4) H. Ashida and A. Kitaoka: A gradient-based model of the peripheral drift illusion. *Perception*, **32** (supplement), 106.
- 5) H. Ouchi: Japanese optical and geometrical art. Dover, Mineola NY, 1977.
- 6) L. Spillmann, F. Heitger and S. Schuller: Apparent displacement and phase unlocking in checkerboard patterns. *Paper presented at the 9th European Conference on Visual Perception*, Bad Nauheim, 1986.
- 7) T. J. Hine, M. Cook and G. T. Rogers: An illusion of relative motion dependent upon spatial frequency and orientation. *Vision Research*, **35**, 3093-3102, 1995.
- 8) T. J. Hine, M. Cook and G. T. Rogers: The Ouchi illusion: An anomaly in the perception of rigid motion for limited spatial frequencies and angles. *Perception and Psychophysics*, **59**, 448-455, 1997.
- 9) B.-G. Khang and E. A. Essock: Apparent swinging motion from a 2-D sinusoidal pattern. *Perception*, **29**, 453-459, 2000.
- 10) B. Pinna and G. J. Brelstaff: A new visual illusion of relative motion. *Vision Research*, **40**, 2091-2096, 2000.
- 11) H. Ashida: Spatial frequency tuning of the Ouchi illusion and its dependence on stimulus size. *Vision Research*, **42**, 1413-1420, 2002.